

## Case Studies



# The use of ochre 40,000 years ago in Africa

---

ARCHAEOLOGY &  
PALEONTOLOGY, CASE STUDY,  
TRIBOLOGY

---

**Alain Queffelec, Daniela Rosso and Francesco d'Errico**



### PACEA | Other articles

PACEA laboratory is a research unit of the Centre National de la Recherche Scientifique (CNRS), the University of Bordeaux, and the French Ministry of Culture. PACEA research mostly focuses on Palaeolithic cultures in Europe and Africa and their environment, biological anthropology, funerary practices, and rock art.

“*Confocal technology is an effective technique to identify facets of ochre pieces ground on different rocks*”

The term “ochre” refers to a variety of rocks characterized by a red or yellow colour or streak, from soil lumps to ore minerals, containing a high proportion of iron oxides. A large number of ochre pieces discovered at this site show traces of utilization, in particular flaking scars and striations produced by grinding. Twenty-one grindstones used for ochre processing were also recovered at the site (Rosso, Pitarch Martí and d’Errico 2016) in the same areas where most of the ochre pieces were found (Rosso, d’Errico and Zilhão 2014).

The purpose of the tribological analysis, conducted in tandem with more classical analytical techniques, was to better understand how ochre pieces were processed and used.

The analyzed samples included ochre pieces bearing facets created by grinding (**Figure 1**). Even though these facets are often described in literature, there have been no attempts to characterize them quantitatively. Experimental grinding of ochre lumps similar to those found at the site on grindstones made of different rocks allowed us to create a comparative framework.

We recorded the variables commonly used to characterize facets (size, presence of grooves or striations, localization on the fragment etc.) and conducted a non-destructive tribological analysis by using a 3D.




**Figure 1. Ochre fragments modified by grinding (top left) and by scraping (bottom left) with close-up views illustrating traces diagnostic of the technique used (modified after Rosso et al. 2017)**

The study was conducted on nineteen archaeological pieces of fine-grained ochre presenting one or more facets, and three very fine-grained, finegrained and coarse-grained experimental ochre pieces. Each of them was abraded on limestone, quartzite and sandstone grindstones.

In order to capture large areas, we measured an area of 5×5 fields of view obtained with the 20X brightfield objective of Sensofar's S neox with a Z-range of few millimetres. Some of the resulting surfaces, after data processing (form and outlier removal, filling of non-measured points) are shown in **(Figure 2)**.






**Figure 2. 3D renderings of experimental (A, B, C) and archaeological (D to L) facets measured using confocal technique (modified after Rosso et al. 2017)**

Gaussian filtering (0.25 mm cut-off) was applied to focus only on roughness and, after testing multiple standard parameters (ISO 25178), Sq and Sdr appeared as the most effective choice to discriminate experimental facets (**Figure 3**). In some cases results highlight significant differences in the roughness values recorded on facets belonging to the same ochre piece.

This suggests that some ochre fragments were processed on different grindstones, arguably at different times, to produce small quantities of ochre powder. Colorimetric and granulometric analyses, conducted in parallel, indicate that the resulting powders differed in colour and grain size and were used for different purposes, either functional or symbolic in nature (Rosso, d'Errico and Queffelec 2017).



**Figure 3. Scatter plots correlating Sq and Sdr values on experimental and archaeological facets. (a) results obtained when grinding very fine-grained ochre (EXP1), fine-grained ochre (EXP2) and coarse-grained ochre (EXP3) on grindstone made of different rocks. (b) Results of the roughness analysis of ground facets on archaeological pieces. Dots of the same colour identify measurements taken on the same facet. Grey dots identify the overall variability of the archaeological sample (modified after Rosso et al. 2017)**

Confocal technology is an effective technique to identify facets of ochre pieces ground on different rocks. Results were instrumental to improving our understanding of ochre processing and use by the Middle Stone Age hominins that inhabited Porc-Epic Cave 40,000 years ago.

Application of the same methodology to collections of ochre pieces from other Middle Stone Age sites may identify changes through time in the way ochre was modified, provide key information on the function it fulfilled in those societies, and help establish when pigments were first used symbolically in the history of mankind.



The use of ochre 40,000 years ago in Africa

## Related products





## More interesting cases



### Prehistoric tool use and the evolution of landscapes: a 3D microwear perspective

ARCHAEOLOGY & PALEONTOLOGY, CASE STUDY, TRIBOLOGY

› Read More



### The use of ochre 40,000 years ago in Africa

ARCHAEOLOGY & PALEONTOLOGY, CASE STUDY, TRIBOLOGY

› Read More

HOME / NEWS / THE USE OF OCHRE 40,000 YEARS AGO IN AFRICA